

Atmospheric Dynamics of Short-period Extra Solar Gas Giant Planets I:

Dependence of Night-Side Temperature on Opacity

Ian Dobbs-Dixon¹ & D.N.C. Lin^{1,2}

¹*Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064, USA*

²*Kavli Institute of Astronomy & Astrophysics, Peking University, Beijing, China*

ABSTRACT

More than two dozen short-period Jupiter-mass gas giant planets have been discovered around nearby solar-type stars in recent years, several of which undergo transits, making them ideal for the detection and characterization of their atmospheres. Here we adopt a three-dimensional radiative hydrodynamical numerical scheme to simulate atmospheric circulation on close-in gas giant planets. In contrast to the conventional GCM and shallow water algorithms, this method does not assume quasi hydrostatic equilibrium and it approximates radiation transfer from optically thin to thick regions with flux-limited diffusion. In the first paper of this series, we consider synchronously-spinning gas giants. We show that a full three-dimensional treatment, coupled with rotationally modified flows and an accurate treatment of radiation, yields a clear temperature transition at the terminator. Based on a series of numerical simulations with varying opacities, we show that the night-side temperature is a strong indicator of the opacity of the planetary atmosphere. Planetary atmospheres that maintain large, interstellar opacities will exhibit large day-night temperature differences, while planets with reduced atmospheric opacities due to extensive grain growth and sedimentation will exhibit much more uniform temperatures throughout their photosphere's. In addition to numerical results, we present a four-zone analytic approximation to explain this dependence.

1. Introduction

Among the over 200 extra-solar planets discovered through a variety of observational techniques including Doppler shifted spectral lines, transit searches, and gravitational lensing, approximately 9% have orbital periods less than 4 days. One of the more puzzling and

unexpected results to come from planet detections, these planets, known as hot-Jupiter's or Pegasus planets, receive approximately 10^4 times more energy from stellar irradiation than from internal heating. This energy input greatly modifies the thermal structure and dynamics of the atmosphere. Given the short orbital period of these planets, tidal forces also play a strong role, quickly circularizing their orbits and synchronizing their spin and orbital frequencies (Goldreich & Soter 1966; Dobbs-Dixon et al. 2004). Consequently, one side of the planet perpetually faces the host star, reaching equilibrium temperatures of over 1000K, while night-side temperatures are determined by the ability of the planet to redistribute energy. In the absence of heat redistribution, the night-side of the planet would remain at $\approx 100\text{K}$ after a few Gyr of cooling and Kelvin-Helmholtz contraction. For comparison, Jupiter has a rotational period of 11 hours and its internal luminosity is approximately equal to the energy it receives from the Sun. Its photospheric temperature is quite uniform in both longitude and latitude.

The small semi-major axis of hot-Jupiter's provide several new methods for exploring their structural parameters and atmospheres including primary transits (Charbonneau et al. 2007), secondary eclipses (Deming et al. 2005b; Charbonneau et al. 2005; Deming et al. 2006), and spectroscopy (Charbonneau et al. 2002; Brown et al. 2002; Deming et al. 2005a; Grillmair et al. 2007; Richardson et al. 2007; Swain et al. 2007). Detection of a primary eclipse, as the planet passes between the star and the Earth, yields measurements of planetary radii and masses that can be used to infer structural parameters of the planets. The secondary eclipse, caused by the passage of the planet behind its host star, provides a direct measurement of the day-side temperature of the objects by measuring the decrement in infrared flux. As of the writing of this paper, 14 planets have been detected via the primary transit method, while 3 of those have also been detected via their secondary eclipse. The day-side temperatures of the objects found from secondary eclipse measurements are found to be $T = 1130 \pm 150\text{K}$ for HD209458b (Deming et al. 2005b), $T = 1060 \pm 50\text{K}$ for TrES-1 (Charbonneau et al. 2005), and $T = 1117 \pm 42\text{K}$ for HD189733b (Deming et al. 2006). Measurements of HD209458b and HD189733b were done using the Spitzer Space Telescopes $24\mu\text{m}$ and $16\mu\text{m}$ band-passes respectively. The estimate of temperature is then sensitive to both the ratio of stellar and planetary radii and stellar temperature. TrES-1 was observed in both the $4.5\mu\text{m}$ and $8\mu\text{m}$ bands, and temperatures were derived by assuming the planet emits as a blackbody. In addition, recent non-transiting observations of ν -Andromeda b with the Spitzer Space Telescope, indicate a substantial orbital phase dependence in the flux at $24\mu\text{m}$ (Harrington et al. 2006).

A number of groups have used one-dimensional, plane-parallel radiative transfer models to calculate the emergent spectra of these hot-Jupiter's (Fortney et al. 2005; Burrows et al. 2005; Seager et al. 2005; Barman et al. 2005; Fortney et al. 2006b; Burrows et al. 2006b; Fortney et al. 2006a). These approaches simultaneously solve the equations of radiative

transfer, radiative equilibrium, and hydrostatic balance to determine the pressure, temperature, and spectral energy distribution as a function of radial position in the atmosphere. The input parameters include the temperatures at the bottom of the atmosphere and the incident stellar energy spectrum at the top. The temperature at the bottom is determined primarily by the internal luminosity, and ideally it should be taken from one-dimensional evolutionary and structural models. However, given the dominate influence of the incident energy near the surface, the choice of the bottom does not noticeably influence the emergent spectra.

In hot-Jupiter's, the intense irradiation of the day-side drives strong thermal winds toward the night-side. The resulting temperature in the upper atmosphere depends on the ability of these winds to redistribute the stellar irradiation, and in general should be a function of both longitude and latitude. In the absence of dynamical models, this redistribution of incident energy at the upper layers is usually set to be some fraction of the incident irradiation, to represent the degree of energy re-distribution. The value of this parameter is a major uncertainty in these radiative models, and authors have computed a number of cases with varying degrees of re-distribution. In addition, such parameterization neglects the role of advection and radiative transfer within lower levels of the atmosphere, which are also important in determining the final pressure-temperature profiles.

In an attempt to address the dynamical redistribution of incident stellar energy within the atmosphere, there have been a number of dynamical models of the atmospheres of hot-Jupiter's. These models utilize a variety of methods employing various simplifications, including solving the primitive equations (Showman & Guillot 2002; Cooper & Showman 2005), the shallow water equations (Cho et al. 2003; Menou et al. 2003; Langton & Laughlin 2007), the equivalent-barotropic equations (Cho et al. 2006), and two-dimensional hydrodynamical equations (Burkert et al. 2005). These models predict a wide range of behavior. More details on these models are presented in §4, in comparison with the model presented in this paper. One group (Fortney et al. 2006a) has attempted to couple the spectral and dynamical models by utilizing pressure and temperature profiles derived directly from the simulations of Cooper & Showman (2005). Unlike previous models, these atmospheres are not iterated to achieve radiative-convective equilibrium.

In this paper, we present the results obtained from a three- dimensional, hydrodynamic simulation based on flux-limited radiative transfer models of hot-Jupiter's for a variety of rotation rates and opacities. In §2, we present the basic equations, numerical methods, and initial conditions. In §3 we present our results and analysis for both rotating and non-rotating flows. We also study the effects of changing opacity on the dynamics and heat distribution. In §4 we include a detailed comparison with previous dynamical models in an attempt to highlight the consequences of making certain simplifying assumptions. We conclude in §5

with a discussion.

2. Numerical Method

2.1. Flux-Limited Radiative Hydrodynamical Model

We cast our numerical model in spherical coordinates (r, ϕ, θ) , where ϕ is the azimuthal angle (or longitude) and θ is the meridional angle (or latitude) measured from the equator. Including both the Coriolis ($2\mathbf{\Omega} \times \mathbf{u}$) and centrifugal forces ($\mathbf{\Omega} \times (\mathbf{\Omega} \times \mathbf{r})$), the equations of motion for the fluid can be written as

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0 \quad (1)$$

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho} \nabla P + \frac{1}{\rho} \nabla \Phi - 2\mathbf{\Omega} \times \mathbf{u} - \mathbf{\Omega} \times (\mathbf{\Omega} \times \mathbf{r}) \quad (2)$$

The rotation frequency is given by Ω , and the gravitational potential, $\Phi = -\frac{GM_p}{r}$, varies only in the radial direction. We neglect explicit viscosity, but some degree of numerical viscosity is inevitable. We also include the curvature terms in $(\mathbf{u} \cdot \nabla) \mathbf{u}$. We solve equations (1) and (2) on a staggered grid, where scalars are defined in the center of cells and vectors on cell boundaries. This method yields second-order spatial accuracy. Given the decreasing grid size near the pole, our computational domain is limited to $\theta = \pm 70^\circ$. Although excluding this region neglects an avenue for energy re-distribution, even a modest amount of rotation (approximately 2 to 4 days for the hot-Jupiter's) will cause the dominate flow patterns to be concentrated near the equator. We simulate the entire azimuth of the planet, instituting periodic boundary conditions at $\phi = 0$ and 2π . The radial extent of the domain extends from $7.95 \times 10^9 \text{ cm}$ to $8.65 \times 10^9 \text{ cm}$, corresponding to 1.06 to $1.2 R_J$. The pressure scale height on the day-side is approximately 340 km.

Our numerical radiative transfer scheme is capable of following the temperature and radiation energy independently, linking them with a given heating/cooling function. However, this degree of sophistication is not necessary for calculations in which the gas is close to thermal equilibrium. Instead, we use a one-fluid approximation where radiation energy density is a simple function of temperature, $E = aT^4$. For the temperatures considered here, the radiative energy density is much lower than the thermal energy density. The internal and radiation energy equations reduce to

$$\left[\frac{\partial \epsilon}{\partial t} + (\mathbf{u} \cdot \nabla) \epsilon \right] = -P \nabla \cdot \mathbf{u} - \nabla \cdot \mathbf{F} \quad (3)$$

where $\epsilon = c_v \rho T$ is the internal energy density, T is the temperature, c_v is the specific heat, and $\mathbf{F}$ is the radiative flux.

To calculate the radiative flux we use the flux limited radiation transfer approximation of Levermore & Pomraning (1981),

$$\mathbf{F} = -\lambda \frac{c}{\rho \kappa} \nabla E, \quad (4)$$

where λ is a non-constant flux limiter which prescribes the relationship between the radiative flux and the radiative energy gradient. We use the flux limiter developed by Levermore & Pomraning (1981), given by

$$\lambda = \frac{2 + R}{6 + 3R + R^2}, \quad (5)$$

where

$$R = \frac{1}{\rho \kappa} \frac{|\nabla E|}{E} \quad (6)$$

compares the scale height of the radiation energy to its mean free path. This overall procedure allows for an accurate treatment of both optically thick and thin regions; in the optically thick *diffusion* limit $R \rightarrow 0$, $\lambda \rightarrow 1/3$, and $\mathbf{F} \rightarrow -c \nabla E / 3 \rho \kappa$, while in the optically thin *streaming* limit $R \rightarrow \infty$, $\lambda \rightarrow 1/R$, and $\mathbf{F} \rightarrow cE$. While the flux limiter λ is an approximation, it is quite accurate in both the optically thick and thin limits.

In order to follow the evolution on a long time-scales, the radiative portion of the energy equation is solved *implicitly*. For quasi-static radiative conditions such as those considered here, the equations can be advanced much more rapidly than if they were restricted by a radiative time-step. We use the successive over-relaxation method (SOR) to solve the $\nabla \cdot \mathbf{F}$ portion of equation (3), alternately updating even and odd grid cells. The limiting factor for the numerical time-step then becomes the Courant condition.

The advection scheme, described in Hawley et al. (1984) and Kley & Hensler (1987), is an extension of the first-order van Leer (van Leer 1977) method known as the 'mono-scheme'. It employs operator-splitting, where the finite difference equations are split into parts, which are then solved separately always using the latest values of the variables. The scheme yields semi-second order temporal accuracy and allows for the accurate resolution of discontinuities in the fluid flow with limited diffusion.

Finally, we use an ideal gas equation of state for the pressure, $p = (R_G \rho T) / \mu$, with specific heat $c_v = R_G / (\mu(\gamma - 1))$ where R_G is the gas constant, $\gamma = \frac{7}{5}$, and the mean molecular weight is fixed at $\mu = 2.3$. Although the temperatures in the hydrodynamic models do become hot enough to dissociate hydrogen molecules in some locations within the planet, the region which we are most concerned with is well described by a constant molecular

weight. Radiative opacities are found using the tables of Pollack et al. (1985) for lower temperatures coupled with Alexander & Ferguson (1994) for higher temperatures. These are Rosseland mean opacities and include the effects of atomic, molecular, and solid particulate absorbers and scatters. The opacity is one of the largest uncertainties. The effect of composition, clouds, settling, wavelength dependence, grains' condensation, sublimation, collisional growth, and sedimentation are but a few of the parameters that alter the magnitude of the opacity. To address this uncertainty, we explore the effect of varying opacity in §3.3.

Our initial density and temperature profiles are taken from a one-dimensional planetary evolutionary code. For details on the model, see model B1 of Bodenheimer et al. (2001). Use of this model implies that the simulations are initially spherically symmetric. Although unrealistic, we allow simulations to run for sufficient time to relax into equilibrium configurations; the details of the initial state are lost. Figure (1) shows the initial temperature and density profiles from the one-dimensional model near the top of the planet. The model follows a $0.63M_J$ planet for 4.5Gyr. The upper boundary was held at a fixed temperature of 1200K to simulate the effects of irradiation from the central star. As noted in Cho et al. (2006), a quiescent initial start neglects the effects of pre-established small-scale structures such as eddies and jets. Given that we lack the resources to simultaneously study these small scale effects and full three dimensional effects, we assume that the dynamics will be dominated by the large scale anisotropic heating imposed by the stellar irradiation. In contrast to the shallow-water approximation, specific vorticity can be generated in our full 3D radiative hydrodynamic simulations. Therefore, baroclinic instabilities can be excited spontaneously and can lead to the generation of structure down to the resolution length scales.

To represent the impinging radiation, we set the temperature at the upper boundary to $T = \max \left[1200\text{K} (\cos(\phi) \cos(\theta))^{1/4}, 100\text{K} \right]$, approximating the stellar radiation field with a maximum at the sub-stellar point, and a night-side held at 100K. The night-side temperature is chosen to be consistent with the photospheric temperature derived in non-irradiated planetary evolution models. This imposed temperature distribution will increase the scale height of the atmosphere on the day-side, while cooling and contracting it on the night-side. An interesting result is a planet that is no longer spherical; the scale-height on the day-side is somewhat larger than on the night-side. Given the computational difficulties of modeling the extremely low density regions in the upper atmosphere, we impose a movable upper boundary at the location where $\rho < 10^{-9}\text{g/cm}^3$. At the bottom boundary we specify the temperature flux taken from our initial one-dimensional models. As mentioned above, the energy input from the star overwhelms the planets intrinsic luminosity in the upper atmosphere, so the choice of this boundary condition is not critical. We have run several simulations with fixed temperature to verify the validity of this approximation, and found little difference in the resulting energy distribution or dynamics.




Fig. 1.— Spherically symmetric temperature and density profiles from the one-dimensional evolutionary models of Bodenheimer et al. (2001) used as an initial condition for the simulations presented here. The models were run for 4.5Gyr, during which time the temperature at the upper boundary was held at 1200K to simulate the irradiation from the central star.

3. Results of Numerical Simulations

3.1. The Non-Rotating Model

As a reference, we first present the results from an idealized and artificial non-rotating model in which the gas giant is subject to one-sided stellar irradiation. The azimuthal pressure gradient associated with the imposed temperature contrast drives strong winds toward the night-side of the planet. The flow pattern is symmetric with respect to the planet-star line, and fluid flows to the night-side around both sides of the planet. As the material moves to the night-side, it achieves maximum velocities of $\sim 3\text{km/s}$ near the terminators ($\phi = \frac{\pi}{2}$ and $\phi = \frac{3\pi}{2}$). This velocity corresponds to Mach numbers of up to 2.7. A substantial portion of energy is released in these shocked regions. By the time the two symmetric flows converge on the night-side they have both undergone substantial cooling and sink radially inward, initiating a return flow at depth. In Figure (2) we show the temperature distribution in the equatorial plane. Despite the high winds, a clear day/night difference persists. Also evident is the cool region at depth, which is due to the confluence of the negative temperature gradient from the interior, the positive temperature gradient near the surface, and the cool flow returning to the day-side, completing the two, approximately symmetric, azimuthal convection cells.

Figure (3) shows the temperature and velocity at the photosphere ($\tau = \frac{2}{3}$) of the planet. Given the changing scale height near the surface of the planet, these plots do not represent a constant radial surface. Moving from the day to night-side, a clear increase in the velocity near the terminator is evident, followed by a drop at the convergent point. It should also be noted that the convergence point is quite dynamical, oscillating in both longitude and latitude. However, despite these oscillations, two distinct convective cells remain. Night-side temperatures range from 300 – 550K, with an average night-side temperature ($\frac{\pi}{2} < \phi < \frac{3\pi}{2}$) of 414K. Also evident in Figures (2) and (3) is an area of increased temperature near 180° . As the two converging flows meet on the night-side, compressional heating drives up the temperatures, and thus the local scale height. The resulting increase is small, but can be seen both at depth and near the photosphere.

3.2. A Planetary Model with 3-Day Rotation

It is widely believed that tidal forces within the atmospheres and the envelope of hot-Jupiter’s drive them to tidally locked spin configurations on time-scales much shorter than the main sequence life span of their host stars. In this synchronous state, the planets’ spin frequency equals their orbital frequency. With this assumption, hot-Jupiter’s spin with periods on the order of 3–days. In comparison to our own giant planets, this is a relatively




Fig. 2.— The temperature distribution at the equator of a non-rotating hot-Jupiter. The inner boundary is assumed to be spherically symmetric, with an outward energy flux fixed from the initial one-dimensional model. The outer black area represents regions with $\rho < 10^{-9} \text{ g/cm}^3$, outside our movable boundary.




Fig. 3.— The temperature (upper panel) and velocity (lower panel) at the photosphere of a non-rotating planet. A strong temperature gradient between the day-side ($\phi = 0^\circ$) and the night-side ($\phi = 180^\circ$) is evident. The regions of largest ∇T correspond to fluid motions of $> 3 \text{ km/s}$.

slow spin rate. Nevertheless, the associated Coriolis force significantly alters the resulting flow dynamics and may have implications concerning the ability of the planet to fully synchronize its spin. Another crucial consideration that came to light during this study was the effects associated with the initialization of the rotation. The one-dimensional initial models described in Section (2.1) were non-rotating models, and the rate that we chose to turn on the rotation had observable effects. We will explore this effect in a subsequent paper.

3.2.1. *The Day-Side Isothermal Surface*

In Figure (4) we show the temperature distribution in the equatorial plane for a simulation rotating with a period of 3-days. The sub-solar point on the day-side is characterized by an radially-extended nearly isothermal region with an effective day-side temperature $T_d \sim 1200K$. Upon adjusting to a hydrostatic equilibrium, a slightly negative temperature gradient is established so that the reprocessed stellar radiation can penetrate into the planetary envelope. Nevertheless the day-side photosphere is essentially isothermal with a density profile

$$\rho(r) = \rho(r_b) \exp [-(g\mu/R_G T_d)(r - r_b)], \quad (7)$$

where r_b is a planet's radius at the base of the isothermal region and $g = GM_p/r_b^2$ is the surface gravity of the planet. The imposed stellar heating falls off as a function of longitude, thus the radial extent of this isothermal region decreases with the inclination angle between the local zenith and the position of the host star overhead.

3.2.2. *Azimuthal Effective-Temperature Distribution*

Below the isothermal photosphere, a cooler region at lower depths is also evident on the day-side, associated in part with a cool return flow from the night-side. Figure (5) shows temperature distribution both throughout the entire photosphere, and focusing on structure on the night-side. Despite the added effect of rotation, a clear day-night delineation is still apparent, with the night-side characterized by effective temperatures T_n from 310 to 500K. The average night-side temperature is 380K, slightly smaller than the non-rotating simulation with the same opacities. This slight decrease in average temperature is due to increased cooling associated with rotationally modified flows discussed in the next subsection. Slightly hotter regions near the terminators, associated with jets from the day-side, are apparent with temperatures reaching $\sim 500K$.




Fig. 4.— The temperature distribution at the equator of a planet rotating with a period of 3days. The day-side is characterized by a large isothermal area near the top, the extent of which falls off with increasing longitude. A cool region at depth is also evident due to the combined effects of a negative temperature gradient from the internal heating, a positive temperature gradient from the irradiation, and a cooling return flow from the night-side.




Fig. 5.— The temperature at the photosphere of a planet rotating with a period of 3days. The upper panel shows the distribution over the entire planet, while the lower panel highlights the temperature structure on the night-side from $\phi = \frac{\pi}{2}$ to $\phi = \frac{3\pi}{2}$. A clear day-night delineation persists, despite complicated dynamical structure, due to substantial radiation near the terminators.

3.2.3. Thermal Current and the Coriolis Effect

The upper panel of Figure (6) shows the velocity magnitude at the photosphere ($|v| = \sqrt{v_\phi^2 + v_\theta^2}$). As in the non-rotating model, material is moving quite rapidly, reaching speeds of $\sim 4\text{km/s}$ near the terminators. Eastward (prograde or $v_\phi > 0$, *i.e.* in the same direction as the unperturbed spin) moving material appears to be funneled from the sub-tropical latitudes ($|\theta| > 0$) into an equatorial jet near $\phi = \frac{\pi}{2}$, while westward (retrograde or $v_\phi < 0$) moving material is pushed from the subtropical zone toward the poles near $\phi = \frac{3\pi}{2}$. It is these flow structures, rapidly advecting energy from the day-side, that account for the hotter regions seen on the night-side in Figure (4). To understand this structure, we must evaluate the $\hat{\theta}$ component of the Coriolis force, given by $-2\Omega v_\phi \sin(\theta)$, which is shown in the lower panel of Figure (6). Although material near the equator doesn't feel any Coriolis force in this direction, it is clear that material at higher and lower (sub-tropical) latitudes does. The asymmetry imposed by the rotation (*i.e.* the azimuthal velocities) causes the fluid moving in eastward and westward directions to behave significantly different than in the non-rotating case.

The approximate magnitude of velocity can be estimated from equation (2), considering only the pressure gradient term. Assuming an approximately constant acceleration around to the night-side of the planet given by $a \simeq -\frac{1}{\rho}\nabla P$, the velocity at the terminator should be given by

$$v_T \simeq \left[\frac{2\gamma R_G}{\mu} (T_d - T_n) \right]^{1/2}. \quad (8)$$

Given a day-side temperature of $T_d \simeq 1200\text{K}$ and an average night-side temperature of $T_n \simeq 350\text{K}$, flows should achieve Mach numbers of ~ 2 near the terminator. The sound-speed $(\gamma R_G T / \mu)^{1/2} \sim 1.7\text{ km/s}$ at the terminator, yielding a local Mach number, as predicted, of ~ 2 .

Also evident in the plot of velocity magnitude in Figure (6) is the marked decrease in velocity where the eastward and westward flows converge. Neither flow is able to instigate circumplanetary flow at the surface. Figure (7) shows an equatorial slice of the azimuthal velocity v_ϕ at the equator (upper panel) and at higher latitudes (lower panel). It is evident from this plot that the eastward moving flow does continue around the planet at depth near the equator, while the westward moving fluid continues around the planet at higher (and lower) latitudes. Because of the effects of rotation shown in Figure (6), the convergence point is near $\phi = \frac{5\pi}{4}$ for the equatorial flow, and near $\phi = \frac{3\pi}{4}$ for flows at higher and lower latitude. This flow pattern implies that, upon converging, one of the two flows has undergone substantially more cooling than its counterpart. Thus, at the equator, when the eastward flow encounters the westward flow near $\phi = \frac{5\pi}{4}$ (past “mid night”), the former is cooler and sinks below. The opposite is true at higher latitudes, with the westward flow experiencing

more cooling and sinking below the eastward flow. This cooling trend can be seen in the lower panel of Figure (5). Lastly, very little motion is apparent deeper in the planet, as the flows are confined to a relatively small region near the top of the planet, supporting our assumption of a spherically symmetric inner boundary condition.

3.2.4. Sub-surface Thermal Stratification

Another method of visualizing the structure of the planet is through pressure-temperature profiles. Of crucial importance when calculating the emergent spectra, the pressure-temperatures at four different longitudes are shown in Figure (8) for both the equator and $\theta = 35^\circ$. All profiles agree well at pressures above 0.1bar, again supporting the assumption of spherical symmetry at depth. However, below this pressure (or above this height), their behaviors are quite different.

The day-side profile undergoes a significant temperature inversion (*i.e.* temperature begins to increase with radius and decrease with pressure). This temperature gradient allows radiative diffusion from the photosphere to the planetary interior. This excess flux is advected to the night-side deep down in the planetary envelope. Comparing to Figure (4), it is evident that the lowest temperature region, $\sim 650\text{K}$, is associated in part with the cool, return flow. This transitional region is analogous to the thermocline in the terrestrial ocean which separates the surface and deep water layers.

Near the upper atmosphere (where the pressure is low), the temperature distribution contains a large, approximately isothermal region on the day-side, clearly dominated by the stellar irradiation. In contrast, the temperature of the night-side ($\phi = \pi$) monotonically decreases throughout the entire atmosphere, reaching a temperature of $\sim 300\text{K}$ at the photosphere. While the day-side is fully radiative, there is a region near the photosphere on the night-side that is convectively unstable in the radial direction. As a consequence of efficient convective transport, the $P - T$ distribution on the night side is approximately adiabatic.

The profiles near the terminators (at both $\phi = \frac{\pi}{2}$ and $\phi = \frac{3\pi}{2}$) exhibit more complex structures due to considerable differences in the advective transport of heat. They also show differences between the equatorial values (left-hand panel) and higher latitudes (right-hand panel). At the equator, the temperature at $\phi = \frac{\pi}{2}$ exhibits an isothermal region ranging from 0.1 to 10^{-3} bars. This terminator is associated with the prograde flow. By the time this flow reaches $\phi = \frac{3\pi}{2}$ it has cooled substantially. A slightly hotter region is evident from the westward flow at lower pressures (larger radius) near the photosphere. For the profiles from $\theta = 35^\circ$, it is at $\phi = \frac{3\pi}{2}$ where an approximately isothermal region exists at depth. The $\phi = \frac{\pi}{2}$ profile at high latitudes decreases monotonically, as very little heat is advected




Fig. 6.— The upper panel shows the magnitude of the velocity at the photosphere, given by $|v| = \sqrt{v_\phi^2 + v_\theta^2}$. The lower panel shows the latitudinal component of the Coriolis effect ($-2\Omega v_\phi \sin(\theta)$) at the photosphere of a planet spinning at 3days. The direction of the latitudinal Coriolis force is different for eastward and westward moving material, causing eastward flows to be focused toward the equator, while westward flow is funneled toward the poles.




Fig. 7.— The velocity distribution at the equator (upper panel) and a latitude of $\theta = 35^\circ$ (lower panel) of a planet rotating with a period of 3days. At the equator, eastward flow (red) is able to circumnavigate the planet at depth, while at higher and lower latitudes, it is westward flow (blue) that is able to traverse all the way around the planet. The sinking of one flow under the other is due to different degrees of cooling due to rotationally altered dynamics.

eastward.

3.3. Opacity Effects

The effect of opacity can not be understated. Opacities regulate the efficiency of both the absorption of the incident stellar irradiation on the day-side and the re-radiation from the night-side. As noted by previous authors, major uncertainties in composition, metallicity, and chemistry all cause significant changes to the opacity. Furthermore, the inclusion of clouds, characterized by models of particle size distributions and vertical extent, tends to smooth out wavelength dependent opacities, resulting in spectral energy distributions that more closely approximate blackbodies. In our current models, opacities are found using the tables of Pollack et al. (1985) for lower temperatures coupled with Alexander & Ferguson (1994) for higher temperatures. These are Rosseland mean opacities and include the effects of atomic, molecular, and solid particulate absorbers and scatters.

In light of the uncertainties associated with opacity, we have studied the effect of uniformly changing the opacity by some multiplicative factor. More detailed studies of specific, temperature and density dependent augmentations to the opacity will be presented elsewhere. Figure (9) shows the temperature distribution across the photosphere of a planet with opacities reduced by both a factor of 100 (upper panel) and a factor of 1000 (lower panel). In comparison to Figure (5), night-side temperatures are significantly higher for both models with lower opacities, and distributions are smoother. In addition, due to changes in the flow detailed below, the hottest spot is displaced slightly from the sub-solar point. At the equator, the displacement is $\sim 10^\circ$ and $\sim 20^\circ$ eastward (in the direction of rotation) for opacity reductions of 100 and 1000 respectively. This displacement is largest at the equator, with maximum temperatures at higher/lower latitudes occurring closer to the sub-solar longitude.

In a previous analysis (Burkert et al. (2005)), we derived a formula for the night-side temperature by equating the radiative timescale with the crossing timescale. This two-zone (day-night) model assumes that: the advective heat flux is larger than the heat flux from the interior, the heat carried by a day-night thermal current is determined by the amount of radiative diffusion during the hemispheric circulation, and the night-side radiates all the heat advected to its proximity as a black body. With these assumptions, the night-side temperature can be estimated by

$$T_n = \left(\frac{4vc_d^2}{3\pi\kappa_d\sigma R_p} \right)^{1/4} \quad (9)$$

which is an decreasing function of opacity. In this formula, v is the average advective speed which is on the order of the day-side sound-speed c_d , and κ_d is the day-side opacity. The




Fig. 8.— The pressure temperature profiles within a planet spinning with a 3day period. The left-hand panel shows profiles at the equator ($\theta = 0$), while the right-hand panel shows profiles from a latitude of $\theta = 35^\circ$. The individual lines represent $\phi = 0$ (solid line), $\phi = \frac{\pi}{2}$ (dotted), $\phi = \pi$ (dashed), and $\phi = \frac{3\pi}{2}$ (dash-dot). The dot on each profile denotes the location of the photosphere.




Fig. 9.— The temperature at the photosphere of a model with the opacities reduced by a factor of 100 (upper panel) and 1000 (lower panel). Lower opacity fluid absorbs the incident stellar irradiation deeper in the atmospheres. This higher density material is able to advect the energy to the night-side more efficiently, leading to larger night-side temperatures.

increase of night-side temperature with decreasing opacity reflects the depth that incident stellar irradiation is deposited on the day-side. If the atmosphere contains grains with an abundance and size distribution comparable to that of the interstellar medium, only shallow heating occurs on the day-side, and the circulation does not effectively transmit heat to the night-side, which then cools well below the day-side. However, as the abundance of grains in the atmosphere is reduced, the stellar radiative flux penetrates more deeply into the atmosphere on the day-side, and the higher density atmospheric circulation carries a larger flux of heat over to the night-side.

For the parameters used here, equation (9) predicts $T_n \sim 250\text{K}$ for the interstellar opacity simulation, while for the lower opacity simulations, the formula predicts $\sim 750\text{K}$ and $\sim 950\text{K}$ for reductions of 100 and 1000 respectively. Inspection of Figures (5) and Figure (9) show average night-side temperatures of $\sim 380\text{K}$, $\sim 700\text{K}$, and $\sim 890\text{K}$ for the same three cases. These results clearly indicate that the night-side temperature decreases with the magnitude of the opacity. The predicted values from equation (9) generally agree with those from the simulation and it provides a framework for understanding the global heat flow. The differences in these results can be attributed to several factors; the surface heat flux carried by the thermal current from the day to night-side is not entirely radiated on the night-side, but rather cools as it travels and advection at depth plays an important role in transporting heat. These effects can be incorporated into a more comprehensive four-zone (day-night and interior-photosphere) model where we examine the energy transfer within the optically thick regions below the planetary photosphere.

Before the presentation of the four-zone model, it is useful to analyze the results of the numerical simulation. These calculations indicate that changing the opacity of the atmosphere not only alters the night-side temperature, but it also modifies both the flow dynamics and interior structure. For large opacity (our standard case), the isothermal region on the day-side is relatively shallow, and thus the increased cooling leads to lower night-side temperatures. The large temperature differential promotes a fast flow velocities around the planet. As the opacities are decreased, and the night-side temperature increases and the velocities decrease. In the lowest opacity simulation, the velocity remains subsonic throughout the entire simulation. A more uniform temperature across the planet surface also allows for circumplanetary flow near the equator by reducing the pressure gradient. Figure (10) shows the velocity at the photosphere of the two low-opacity simulations. These results should be compared to the standard-value results shown in Figure (6). As in the standard case, flow at higher and lower latitudes still travels westward at the $\frac{3\pi}{2}$ terminator and circumplanetary flow at the surface is suppressed due to increased cooling times.

In Figure (11) we show the pressure-temperature profiles at the equator for the two reduced opacity simulations. In contrast to Figure (8) where the night-side was fully con-

vective, both lower opacity simulations exhibit isothermal regions in the upper atmospheres around the entire planet. Below the photosphere, there is a slightly negative temperature gradient so that the day-night advective heat flux deep beneath the photosphere can radiatively diffuse to the planet’s photosphere. As it is to be expected, the radial extent of the nearly isothermal atmosphere is largest with the lowest opacity. Both simulations retain a convective regions below the isothermal regions on the night-side.

3.3.1. *A Four-Zone Model for the Sub-Surface Thermal Structure of Planetary Atmospheres*

Discussions in the previous section clearly indicate that thermal currents not only transport heat from the day to night, but also from the night to day-side at other radii. In addition, the day-side atmosphere is radiative and isothermal, while the night-side is convective and adiabatic. Finally, quasi hydrostatic equilibria is maintained in most regions except the subduction zones which separate the thermal currents. We now take these effects into consideration in the determination of the night-side temperature and the sub-surface thermal structure.

The two-zone approximation in equation (9) is based on the assumption that the radiative and crossing time-scales in the planetary atmosphere are nearly equal. We show in the next section (Figure (12)) that this assumption is only marginally satisfied on the night-side near the photosphere. In the standard opacity model, the radiative timescale around the terminators is several orders of magnitude shorter than the crossing timescale. This allows for significant cooling, yielding much lower night-side temperatures than would be expected from equation (9). Similar behavior is seen in the lower opacity models.

In order to analytically account for the transport of heat well below the photosphere on either side of the planet, we divide it into four zones, representing the day-night and interior-photosphere regions. There are then 4 sets of thermodynamic variables including: P_{DS} , P_{DI} , P_{NS} , P_{NI} , T_{DS} , T_{DI} , T_{NS} , and T_{NI} where P and T are pressure and temperature, and the subscripts D , N , S , and I represent day, night, photospheric surface and interior respectively. There are three additional quantities which connect these regions: the velocity between the day and night interiors, v_{adv} , the radial distance between the photospheric surface and the planetary interior, l_D , and the density of the gas at the planetary interior on the day-side ρ_{DI} . These quantities can be solved simultaneously with the following 11 equations.

Although the total stellar incident flux $F = L_*/4\pi D^2$ (where L_* is the stellar luminosity and D is the distance between the star and the planet), irradiates only on the day-side, the




Fig. 10.— Velocity at the photosphere of models with opacities reduced by a factor of 100 (upper panel) and 1000 (lower panel). Velocities, determined by both the day-night temperature differential and the cooling efficiency, decrease with decreasing opacity.




Fig. 11.— The pressure temperature profiles at the equator of a planet spinning with a 3day period. The left-hand panel shows profiles for the run with opacities reduced by a factor of 100, while the right-hand panel shows profiles for a simulation with opacities reduced by a factor of 1000. The individual lines represent $\phi = 0$ (solid line), $\phi = \frac{\pi}{2}$ (dotted), $\phi = \pi$ (dashed), and $\phi = \frac{3\pi}{2}$ (dash-dot). The dot on each profile denotes the location of the photosphere, considerably deeper then in the standard opacity simulation.

condition for thermal equilibriums implies that

$$2\sigma(T_{DS}^4 + T_{NS}^4) = F. \quad (10)$$

Hydrostatic equilibrium at the photosphere of both the day and night sides gives:

$$P_{DS} = \frac{2}{3} \frac{g}{\kappa_{DS}}, \quad P_{NS} = \frac{2}{3} \frac{g}{\kappa_{NS}}. \quad (11)$$

For the simplicity of analytic approximation, we represent opacity as

$$\kappa = \kappa_0 T^\beta. \quad (12)$$

In the temperature and density ranges which are relevant to the present investigation, the opacity is roughly independent of density, thus we adopt the approximation $\kappa_0 = 0.0391$ and $\beta = 0.641$.

Our numerical results indicate that the pressure may change by more than two orders of magnitude between the planetary surface and the temperature inversion layer. In contrast, the temperature changes by less than a factor of two. In the spirit of analytic simplicity, we adopt an isothermal approximation for the hydrostatic envelope when we determine the pressure at the planetary interior on the day-side, such that

$$P_{DI} = P_{DS} \exp \left[\frac{l_d g}{c_{s,d}^2} \right], \quad (13)$$

where $c_{s,d} = (\gamma R_G T_{DS} / \mu)^{1/2}$ is the sound speed. However, to calculate the temperature variation, we adopt a radiative diffusion approximation in the radial direction. Including the azimuthal advective transport, we have

$$1 - \left(\frac{T_{DI}}{T_{DS}} \right)^{4-\beta} \simeq (4 - \beta) \left[\left(1 - \frac{\rho_{DI} v_{adv} c_{s,d}^2}{F} \right) \exp \left[\frac{g l_d}{c_{s,d}^2} \right] - 1 \right] \left(1 + \frac{T_{NS}^4}{T_{DS}^4} \right), \quad (14)$$

where

$$\rho_{DI} = \mu P_{DI} / R_G T_{DI}. \quad (15)$$

We choose l_D to be the depth where advection carries half of the incident flux, *i.e.* where

$$\rho_{DI} v_{adv} c_{s,d}^2 \simeq F/2. \quad (16)$$

The advective velocity at depth is driven by the pressure differential, such that

$$v_{adv}^2 \simeq \frac{2 R_G T_{DI}}{\mu} \ln \left(\frac{P_{DI}}{P_{NI}} \right). \quad (17)$$

We assume the consequence of the advective transport is to thoroughly mix the gas so that planetary interior becomes isothermal with

$$T_{NI} = T_{DI} \quad (18)$$

For the standard opacity model, the night-side remains fully convective. Assuming convection is efficient, the envelope of the night-side is expected to be adiabatic, such that

$$\frac{T_{NI}}{T_{NS}} = \left(\frac{P_{NI}}{P_{NS}} \right)^{\frac{\gamma-1}{\gamma}}. \quad (19)$$

The energy equation for the night-side can be written as

$$\left(1 - \frac{P_{DI}}{P_{NI}} \right) = - \frac{v_{conv} g \mu \pi R_p}{v_{adv} R_G T_{NI}} \quad (20)$$

where the convective velocity v_{conv} can be obtained from the mixing length theory. For the low opacity models, the upper regions of the night-side become stabilized against convection. We can then replace the energy equation with the diffusion approximation and recalculate the thermal structure accordingly.

With these equations, we can provide a set of algebraic equations which essentially reproduce the behavior of our numerical simulation. These equations are also more comprehensive than that in equation (9). For opacity similar to that of the interstellar medium (*i.e.* no reduction in κ_0), the night-side remains convective so that equations(10)-(20) are valid. In this limit, T_{DS} is sufficiently larger than T_{NS} such that equations (10) and (11) lead to complete information on the surface layer of the day-side,

$$T_{DS}^4 = F/2\sigma, \quad P_{DS} = 2g/3\kappa_0 T_{DS}^\beta. \quad (21)$$

In comparison with the large variations in equation (13), the day-side is nearly isothermal with $T_{DI} \sim T_{DS}$. At the depth where half of the incident stellar flux is advected from the day to night-side, equation (16) yields

$$P_{DI} v_{adv} \simeq \frac{F}{2}, \quad l_d \simeq \frac{c_{s,d}^2}{g} \ln \left(\frac{3F\kappa_0 T_{DS}^\beta}{4g v_{adv}} \right). \quad (22)$$

In the fully adiabatic night-side, we find from equations (11), (13), (17), and (19) that

$$P_{NS} = P_{DS} (T_{NI}/T_{NS})^\beta, \quad (23)$$

and

$$\frac{T_{NS}}{T_{DS}} = \left[\exp \left(\frac{v_{adv}^2 - 2gl}{2c_{s,d}^2} \right) \right]^{(\gamma-1)/(\gamma+\beta(\gamma-1))}. \quad (24)$$

From equations (17), (18), and (20), we find

$$\exp \left[\frac{v_{adv}^2}{2c_{s,d}^2} \right] - \frac{v_{conv}}{v_{adv}} \frac{g\pi R_p}{c_{s,d}^2} = 1. \quad (25)$$

In the limit that $v_{adv} < c_{s,d}$, equation (25) reduces to

$$v_{adv} \sim (2v_{conv}g\pi R_p)^{1/3}. \quad (26)$$

The convective speed can be estimated from the mixing length theory which is mostly determined by $T_{NI} \sim T_{DS}$ and ρ_{NI} . These quantities have very little or no dependence on κ_0 so that v_{adv} does not change significantly as a function of κ_0 . In this limit, equation (24) becomes

$$\frac{T_{NS}}{T_{DS}} \propto \left[\frac{4gv_{adv}}{3F\kappa_0 T_{DS}^\beta} \right]^{((\gamma-1)-1)/(\gamma+\beta(\gamma-1))} \propto \kappa_0^{-1/4}. \quad (27)$$

Thus, the four-zone model generates a similar T_{NS} dependence on κ_0 as equation (9). In comparison with numerical results, equation (27) does reasonably well for high opacity convective simulations, but expression would be improved by considering the possibility that the night-side may also become radiative in the limit of very low κ_0 .

4. Analysis and Model Comparisons

A number of groups, including Showman & Guillot (2002), Cho et al. (2003), Burkert et al. (2005), Cooper & Showman (2005), Cho et al. (2006), and Langton & Laughlin (2007) have carried out non-linear numerical simulations studying the dynamics of hot-Jupiter atmospheres. Both the methods and results vary considerably. In this section, we attempt to compare their methodology to that presented here, concentrating on the assumptions that lead to differing results.

Both Showman & Guillot (2002) and Cooper & Showman (2005) solve the primitive equations, the former using the EPIC code developed by Dowling et al. (1998) and the later using the ARIES/GEOS Dynamical Core model initially developed by Suarez et al. (1995). The primitive equations, cast in isobaric coordinates, are widely used in meteorology. When deriving them from the full Navier-Stokes equation, a key assumption of hydrostatic equilibrium is invoked. Assumption of hydrostatic equilibrium explicitly links the thickness of a layer to its local temperature. Radial motion can only occur in conjunction with divergence along isobars, the magnitude determined by that which maintains the condition of hydrostatic equilibrium. With this assumption, the continuity equation and hydrostatic condition take the place of the radial momentum equation, and can be written as

$$\nabla_p \cdot \mathbf{v} + \frac{\partial \omega}{\partial p} = 0, \quad (28)$$

where $\mathbf{v}$ is the horizontal velocity and ω is the vertical velocity with respect to pressure coordinates. The resulting radial velocity is slow in comparison to the horizontal (or isobaric) motion. Although an excellent assumption for thin terrestrial atmospheres, it neglects critical flows present in thicker atmospheres such as those of hot-Jupiter's. Although deviation from hydrostatic equilibrium provides the dominate radial acceleration, high velocity azimuthal flows on the night-side give rise to non-negligible radial forces neglected in the primitive equations that contribute to the ability of the converging flows to pass under one another as seen in Figure(7).

Our numerical results and analytic approximation clearly indicate the dependence of the night-side temperature on the efficiency of radiation transfer. The heat flux carried by the thermal current is determined by the penetration depth of the incident stellar radiation. Radiative diffusion and convection are essential processes which regulate the heat diffusion well below the photosphere of the planet. In conjunction with the primitive equations, both Showman & Guillot (2002) and Cooper & Showman (2005) utilize a Newtonian radiative scheme to approximate stellar irradiation. In principle, this scheme is only appropriate for energy deposition and emission in optically thick regions. The Newtonian radiative scheme provides a heating/cooling term to the energy equation that relaxes the temperature toward some pre-defined equilibrium distribution on some radiative timescale. The forcing term can be expressed (Cooper & Showman 2005) as,

$$\frac{q}{c_v} = -\frac{T(\theta, \phi, p, t) - T_{eq}(\theta, \phi, p)}{\tau_{eq}(p)}. \quad (29)$$

For the radiative timescale, τ_{eq} , Showman & Guillot (2002) assume a constant value given by that at 5bars, while Cooper & Showman (2005) use the calculations of Iro et al. (2005) to set a radiative relaxation timescale that is dependent on the local pressure. As noted by the authors, this approximation is crude, but allows rapid computation of a large number of models. In order for the Newtonian approximation to be viable, the radiative timescale ($\tau_{rad} \approx \frac{E_T}{F}$) must be much longer than the crossing timescale ($\tau_x \approx \frac{\pi R_p}{2|v|}$). In this limit, the temperature distribution will be determined by the dynamics, rather than the assumed equilibrium distribution. This assumption becomes problematic in the upper atmosphere. In Figure (12) we plot the ratio of timescales $\frac{\tau_{rad}}{\tau_x}$ at the photosphere of the planet.

It is clear from Figure (12) that $\tau_{rad} < \tau_x$ throughout the photosphere. The most significant deviation occurs on the day-side, where there is very little motion, and τ_{rad} exceeds τ_x at the planetary photosphere by many orders of magnitude. Near the terminator, where the stellar irradiation falls drastically, the surface radiative timescale is also much shorter than the dynamical timescale; it is here that the deviations from a Newtonian radiative scheme are most important. The flow radiates a significant portion of its energy before it flows to the night-side, allowing for the sharp edges in the temperature distribution seen




Fig. 12.— The ratio of radiative timescale (τ_{rad}) to the crossing timescale (τ_x) at the photosphere of the simulation with interstellar opacities. Near the terminator, the primary region of interest, the radiative timescale is ~ 5 orders of magnitude shorter than the crossing timescale. This rapid cooling accounts for the sharp temperature gradient near the terminator seen in Figure (5).

in Figure (5), despite high velocity flows. In contrast, the Newtonian assumption that $\tau_{rad} \gg \tau_x$, implies that the fluid carries a significant quantity of heat, leading to an overall distribution that will be more uniform with longitude. Finally, despite substantial radiation near the terminators, the near equality of τ_{rad} and τ_x at $\phi = \pi$ demonstrates that the temperature at the back-side is determined purely by the advection of energy by the flow.

To again highlight the effect of opacity, Figure (13) shows the ratio of radiative and crossing timescales, $\frac{\tau_{rad}}{\tau_x}$, for the runs with varying opacity. The upper and lower panels show the simulations with opacities reduced by a factors of 100 and 1000 respectively. Again, the most important area in determining the behavior is near the terminators. In comparison to Figure (12), it is obvious that lower opacity fluid is able to advect energy to the night-side more efficiently due to increased cooling times. As mentioned above, lower opacity on the day-side, allows heat to penetrate further into the planet, and re-radiation near the terminator is somewhat suppressed. Although Figure (13) shows a marked increase in $\frac{\tau_{rad}}{\tau_x}$ from the interstellar opacity simulation, the Newtonian approximation ($\tau_{rad} \gg \tau_x$) is never satisfied. The largest values, $\frac{\tau_{rad}}{\tau_x} \sim 3$, only occur on the night-side in the high velocity circumplanetary jet.

The studies of Cho et al. (2003), Menou et al. (2003), Cho et al. (2006), and Langton & Laughlin (2007) take a different approach, concentrating on the effect of eddies and waves on the overall dynamics. Cho et al. (2003) Menou et al. (2003) and Langton & Laughlin (2007) use the shallow-water equations, while Cho et al. (2006) solves the more generalized equivalent-barotropic equations. The equivalent-barotropic equations can be derived by vertically integrating the primitive equations described above. The resulting equations describe fluid flow in a single, isentropic layer whose scale height can vary. Stellar heating is prescribed by a deflection of the scale-height as a function of position. Concentrating on a single layer allows for the high resolution necessary to study the effect of turbulent eddies and waves. However, allowing for only a single layer implies that the atmosphere is radial isothermal. Although not a bad assumption in the upper regions on the day-side, the rest of the atmosphere exhibits significant radial temperature structure. As noted above, cooling in the upper regions of the night-side allows material to sink radial and initiate a return flow. Disallowing this flow would significantly alter the resulting dynamics.

The final dynamical study of hot-Jupiter's was done by Burkert et al. (2005). The numerical model they used is quite similar to the model presented in this paper. They solve the full hydrodynamical equations, given by equations (1), (2) and (3) together with flux-limited radiation diffusion. However, they restrict their attention to the $r - \phi$ plane from $\phi = 0 \rightarrow \pi$, neglect the curvature terms in equation (2), and ignore the effects of rotation. Comparing our non-rotating results to Burkert et al. (2005) 'Case 1' with our 'standard' opacity, we find that the two models agree quite well. The only substantial differences




Fig. 13.— The ratio $\frac{\tau_{rad}}{\tau_x}$ at the photosphere of models with opacities reduced by a factor of 100 (upper panel) and 1000 (lower panel). Although lowering the opacity decreases the ability of the fluid to radiate its thermal energy, the radiative timescale remains several orders of magnitude shorter than the crossing timescale for most of the photosphere.

are that our backside temperature is slightly lower ($\sim 415\text{K}$ compared to their 480K), and the convergence point is seen to oscillate in the simulations presented here. Our cooler temperatures can easily be explained by considering the increased compressional heating that the fluid in Burkert et al. (2005) simulations experiences as it hits the boundary at $\phi = \pi$, and our extension into three-dimensions. Flows that are able to spread in latitude will cool more than those confined to the equator. Given that our non-rotating simulation results in two symmetric convective cells suggests that by restricting attention to $\phi = 0 \rightarrow \pi$, they did not miss any fundamental physics for that scenario. However, the addition of rotation significantly changes flow patterns and allows for increased cooling.

In summary, four fundamental differences are included in the models presented here, that are not included in some way in the previous methods: self-consistent radiative transfer, solution of the full radial momentum equation, 3-dimensions, and rotation. Noticeable changes occur due to the inclusion of each. Self-consistent radiative transfer allows significant cooling as the fluid travels around to the terminators and night-side of the planet. Coriolis forces alter the structure of the flows, alternately compressing and diverging streamlines moving around the planet in different directions, and the multi-dimensional aspect changes the resulting temperature distribution. Finally, the full treatment of the radial momentum equation yields significant, non-hydrostatic radial structure and motion. Disallowing efficient cooling, and limiting radial motion, while subjecting the planet to the continual stellar energy input, will lead to much more uniform temperature distributions across the entire planet than those that we observe here.

5. Discussion

In this paper we have considered the flow dynamics and heat redistribution in a tidally locked hot-Jupiter orbiting a solar-type star with an orbital period of 3days . We utilized a three-dimensional model that solves the full hydrodynamical equations and models radiative transfer the flux-limited radiation diffusion. We show that the temperature distribution across the planetary atmosphere is a sensitive function of its opacity. Our models exhibit strong day-night temperature contrasts, despite strong winds, the size of which increases with increasing atmospheric opacity. Large temperature differences are due, in a large part, to significant cooling of the flow near the terminators of the planet. Rotational effect significantly alter the flow pattern, allowing for increased cooling and suppressing surface circumplanetary flow in the higher opacity atmospheres.

In our standard model, opacities are calculated using the tables of Pollack et al. (1985) for lower temperatures coupled with Alexander & Ferguson (1994) for higher temperatures. These are interstellar Rosseland mean opacities and include the effects of atomic, molecular,

and solid particulate absorbers and scatters. Although there is significant room for improvement in our treatment of opacity, we explore the effect of uniformly reducing them by factors of 100 and 1000. In agreement with the two-dimensional simulations of Burkert et al. (2005), we find that lower opacities yield higher night-side temperatures. Lower opacities allow incident stellar irradiation to be deposited at larger depths on the day-side, decreasing the cooling rate as this energy is advected to the night-side, and increasing night-side temperatures. The relation between the night-side effective temperature and the opacity in the planetary atmosphere is verified by a more comprehensive four-zone analytic model which takes detailed account of the effect of radiation transfer on both the day and night-side of the planet. This model highlights the importance of appropriately treating radiation transfer in the simulation of atmospheric dynamics on close-in gas giant planets.

The actual value of opacity in the atmospheres of these planets is highly uncertain. However, the day-side temperature of short period gas giant planets around solar type stars is $T_{DS} \sim 1200\text{K}$, which is below the grain destruction temperature. We note that the flow returning to the day-side is dredged up from a cooler interior ($T_{DI} < T_{DS}$) at the sub-solar point. Thus, throughout the thermal circulation, refractory magnesium-rich grains are preserved. Other species of less refractory silicate grains will sublime near the planetary photosphere on the day-side, condense on night-side surface, and be carried along with the returning current beneath the surface. These thermal currents become turbulent as they generate specific vorticity and excite instabilities. Small grains are well coupled to the gas and turbulence induces them to collide and coagulate. In a static atmosphere, particles with sizes s_p and density ρ_p attain terminal velocities of

$$v_t \sim (g\rho_p s_p / \rho_g)^{1/2} \quad (30)$$

in a background gas with density ρ_g . Since the distance over which these particle attain their terminal speed, $L_{term} \sim \rho_p s_p / \rho_g$, is smaller than the density scale height $\sim c_{s,d}^2 / g$, particles larger than $\sim 1\text{mm}$ cannot be carried by the upwelling current on the day-side, and are left well beneath the planet's photosphere. This potential channel for grain-gas separation implies that it is very likely that the opacity in the planet's atmosphere is well below that of the interstellar medium. A detailed analysis of the grain evolution in the atmosphere of short-period gas giants will be presented elsewhere.

We also presented a detailed comparison of our model to previous dynamical models of Showman & Guillot (2002), Cho et al. (2003), Burkert et al. (2005), Cooper & Showman (2005), Cho et al. (2006), and Langton & Laughlin (2007). The approaches among these groups varies greatly, and we have attempted to highlight several of the crucial differences. The most significant differences that we include are solving the full fluid equations in all three-dimensions, rotation, and our treatment of radiation via the flux-limited diffusion method. Flux-limited radiation transport allowed for a self-consistent treatment of the flow

of radiation throughout the planet. This treatment produced significantly shorter radiative timescales than previously calculated, yielding much lower night-side temperatures.

The dynamics and heat distribution within the atmospheres giant planets allow us to probe fundamental questions surrounding planet formation. The diversity of planetary sizes must ultimately result from interior structure variations arising from formation and evolutionary processes. These atmospheres not only serve as a valuable observable links to the interior, but may also play a role in regulating planetary contraction rates. As suggested in Burrows et al. (2000) and Burrows et al. (2006a) an isothermal atmosphere will result in decreased heat transfer and increased planetary contraction timescales. However, a strong day-night temperature difference may allow an avenue for cooling via the night-side. Based on the simulations presented here, it is clear that the efficiency of redistribution decreases with increasing opacity. High opacity atmospheres, similar to interstellar values, will have cool night-side temperatures, providing an avenue for internal heat loss and radial contraction. If, as is expected, there is a significant amount of grain growth and sedimentation, thus reducing the atmospheric opacity, the planets night-side will also contain a large isothermal region, possibly allowing for the retention of internal energy. Diversity in atmospheric opacity may lead to diversity in planetary radii. With the promise of improved observational techniques, including transit spectroscopy and full phase light-curves, the relation between opacity and temperature differential should be testable in the near future.

We thank Geoff Bryden, Peter Bodenheimer, Katherine Kretke, and Gordon Ogilvie for supplying the original version of our numerical scheme, initial structural model, and many helpful discussions. We would also like to acknowledge the use of NASA's High End Computing Program computer systems. This work is partially supported by NASA (NAGS5-11779, NNG04G-191G, NNG06-GH45G), JPL (1270927), NSF(AST-0507424, PHY99-0794) and IGPP.

REFERENCES

- Alexander, D. R. & Ferguson, J. W. 1994, *ApJ*, 437, 879
- Barman, T. S., Hauschildt, P. H., & Allard, F. 2005, *ApJ*, 632, 1132
- Bodenheimer, P., Lin, D. N. C., & Mardling, R. A. 2001, *ApJ*, 548, 466
- Brown, T. M., Libbrecht, K. G., & Charbonneau, D. 2002, *PASP*, 114, 826
- Burkert, A., Lin, D. N. C., Bodenheimer, P. H., Jones, C. A., & Yorke, H. W. 2005, *ApJ*, 618, 512

- Burrows, A., Guillot, T., Hubbard, W. B., et al. 2000, *ApJ*, 534, L97
- Burrows, A., Hubeny, I., Budaj, J., & Hubbard, W. B. 2006a, *ArXiv Astrophysics e-prints*
- Burrows, A., Hubeny, I., & Sudarsky, D. 2005, *ApJ*, 625, L135
- Burrows, A., Sudarsky, D., & Hubeny, I. 2006b, *ApJ*, 650, 1140
- Charbonneau, D., Allen, L. E., Megeath, S. T., et al. 2005, *ApJ*, 626, 523
- Charbonneau, D., Brown, T. M., Burrows, A., & Laughlin, G. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil, 701–716
- Charbonneau, D., Brown, T. M., Noyes, R. W., & Gilliland, R. L. 2002, *ApJ*, 568, 377
- Cho, J. Y.-K., Menou, K., Hansen, B., & Seager, S. 2006, *ArXiv Astrophysics e-prints*
- Cho, J. Y.-K., Menou, K., Hansen, B. M. S., & Seager, S. 2003, *ApJ*, 587, L117
- Cooper, C. S. & Showman, A. P. 2005, *ApJ*, 629, L45
- Deming, D., Brown, T. M., Charbonneau, D., Harrington, J., & Richardson, L. J. 2005a, *ApJ*, 622, 1149
- Deming, D., Harrington, J., Seager, S., & Richardson, L. J. 2006, *ApJ*, 644, 560, secondary Eclipse Discovery Paper for HD189733b
- Deming, D., Seager, S., Richardson, L. J., & Harrington, J. 2005b, *Nature*, 434, 740
- Dobbs-Dixon, I., Lin, D. N. C., & Mardling, R. A. 2004, *ApJ*, 610, 464
- Dowling, T. E., Fischer, A. S., Gierasch, P. J., et al. 1998, *Icarus*, 132, 221
- Fortney, J. J., Cooper, C. S., Showman, A. P., Marley, M. S., & Freedman, R. S. 2006a, *ApJ*, 652, 746
- Fortney, J. J., Marley, M. S., Lodders, K., Saumon, D., & Freedman, R. 2005, *ApJ*, 627, L69
- Fortney, J. J., Saumon, D., Marley, M. S., Lodders, K., & Freedman, R. S. 2006b, *ApJ*, 642, 495
- Goldreich, P. & Soter, S. 1966, *Icarus*, 5, 375
- Grillmair, C. J., Charbonneau, D., Burrows, A., et al. 2007, *ArXiv Astrophysics e-prints*
- Harrington, J., Hansen, B. M., Luszcz, S. H., et al. 2006, *Science*, 314, 623

- Hawley, J. F., Wilson, J. R., & Smarr, L. L. 1984, *ApJS*, 55, 211
- Iro, N., Bézard, B., & Guillot, T. 2005, *A&A*, 436, 719
- Kley, W. & Hensler, G. 1987, *A&A*, 172, 124
- Langton, J. & Laughlin, G. 2007, *ApJ*, 657, L113
- Levermore, C. D. & Pomraning, G. C. 1981, *ApJ*, 248, 321
- Menou, K., Cho, J. Y.-K., Seager, S., & Hansen, B. M. S. 2003, *ApJ*, 587, L113
- Pollack, J. B., McKay, C. P., & Christofferson, B. M. 1985, *Icarus*, 64, 471
- Richardson, L. J., Deming, D., Horning, K., Seager, S., & Harrington, J. 2007, *ArXiv Astrophysics e-prints*
- Seager, S., Richardson, L. J., Hansen, B. M. S., et al. 2005, *ApJ*, 632, 1122
- Showman, A. P. & Guillot, T. 2002, *A&A*, 385, 166
- Suarez, M. J., Takacs, L. L., Suarez, M. J., & Takacs, L. L. 1995, in *Unknown*
- Swain, M. R., Bouwman, J., Akeson, R., Lawler, S., & Beichman, C. 2007, *ArXiv Astrophysics e-prints*
- van Leer, B. 1977, *Journal of Computational Physics*, 23, 276